

Revive Biote Male Post-Insertion Instructions

- Your insertion site has been covered with two layers of bandages. The inner layer is a steri-strip, and the outer layer is a waterproof dressing.
- Do not take tub baths or get into a hot tub or swimming pool for 7 days. You may shower, but do not remove the steri-strips for 7 days.
- No heavy lifting or major exercises for the incision area for the next 7 days, which includes running, elliptical, squats, lunges, etc.
- The sodium bicarbonate in the anesthetic may cause the site to swell for 1-3 days.
- The insertion site may be uncomfortable for up to 2 to 3 weeks. If there is itching or redness you may take Benadryl for relief (25 to 50 mg orally every 6 hours). Caution: this can cause drowsiness!

- You may experience bruising, swelling, and/or redness of the insertion site which may last from a few days up to 2 to 3 weeks. If the redness worsens after the first 2-3 days, please contact the office.
- You may notice some pinkish or bloody discoloration of the outer bandage. This is normal.
- If you experience bleeding from the incision, apply firm pressure for 5 minutes.
- Please call if you have any bleeding not relieved with pressure (not oozing), as this is NOT normal.
- Please call if you have any pus coming out of the insertion site, as this is NOT normal.
- We recommend putting an ice pack on the area
 where the pellets are located a couple of times for
 about 20 minutes each time over the next 4 to 5
 hours. You can continue this for swelling, if needed.
 Be sure to place something between the ice pack
 and your bandages/skin. Do not place ice packs
 directly on bare skin.

REMINDERS

- Remember to have your post-insertion blood work done 4 weeks after your FIRST insertion. If you are not feeling any better by 4 weeks, however, please call the office to have your labs drawn early.
- Most men will need re-insertion of their pellets 4-5 months after their initial insertion. If you experience symptoms prior to this, please call the office.
- Please call as soon as symptoms that were relieved from the pellets start to return to make an appointment for your next insertion.